



INSTRUCTIONS FOR USE:

RECOMMENDED INDICATIONS:

IOS Orthodontic Glass Ionomer LC Band Cement is a fluoride releasing cement formulated for:

- 1. Cementation of orthodontic bands.
- 2. Cementation of appliances retained with stainless steel crowns

CONTRAINDICATIONS:

In rare cases the product may cause sensitivity in some people. If any such reactions occur, discontinue the use of the product and refer to a physician.

COMPOSITION:

Fluoro-alumino-silicate glass

Dimethacrylate

Silicon dioxide

Photo initiators

Stabilizers

PRODUCT PRESENTATION

5 gm Syringe # 17-211-100

INSTRUCTIONS FOR USE:

TOOTH AND BAND PREPARATION

- 1. After removal of separators, scale gingival margins to remove any plaque.
- 2. Pumice or prophy teeth to be banded. If using prophy paste, use oil-free products only.
- 3. Make sure bands are prepared and handled according to manufacturer's instructions.
- 4. Trial fit bands.
- 5. Clean bands before cementing.

CEMENTATION

- 1. Isolate teeth to be banded.
- 2. Slowly extrude the cement from the syringe. Using a small spatula or similar tool, coat the internal surface of the band with a thin layer of the band cement
- 3. While performing this procedure, shield the cement from direct exposure to intense light to avoid premature curing of cement.
- 4. Immediately after using cement, put the cap on the syringe.
- 5. Teeth must be dry prior to final seating of bands.
- 6. Firmly seat the band.
- 8. If using conventional curing light, cure the cement from the occlusal side for 40-50 seconds.
- 9. Refer to light manufacturer's guide for complete instructions.

WARNING:

Avoid looking directly at the curing light when curing Band Guard Cement. Wear protective glasses

USE AND STORAGE:









Store away from sunlight

Observe storage temperatures

Protect from moisture

Shelf life, 3 years

EMERGENCY MEASURES:

Eye Contact: Flush with copious amounts of water. Seek medical attention if irritation persists.

Ingestion: No harm anticipated; if large quantities are swallowed seek medical attention. Advise of fluoride presence in material.

Skin Contact: Wash with soap and water.

Inhalation: No harm anticipated.

Caution: Federal law restricts this device to sale by or on the order of a dentist (or dental professional).

International Orthodontic Services
12811 Capricorn St. Stafford, TX 77477 USA.
T: +1 (832) 342 9487 F: +1 (281) 340 4794
www.iosortho.com info@iosortho.com