

GLASS IONOMER CEMENT

INSTRUCTIONS FOR USE:

RECOMMENDED INDICATIONS:

IOS Glass Ionomer Cement represents an advanced, fluoride-releasing glass ionomer formulation designed for a broad scope of uses:

- 1. Cementation of metal or porcelain fused to metal crowns, bridges, inlays.
- 2. Cementation of stainless steel crowns or cementation of orthodontic bands.
- 3. Base or liner.

CONTRAINDICATIONS:

Pulp capping and sensitive teeth.

COMPOSITION:

Alumino silicate glass Polyacrylic acid

PRODUCT PRESENTATION:

Regular Kit (15gm Powder/15ml Liquid) # 17-211-102 Economy Kit (100gm Powder/60ml Liquid) # 17-211-103

INSTRUCTIONS FOR USE:

- 1. Prepare tooth cavity and base as required.
- 2. Clean and dry enamel and dentin thoroughly.
- 3. Shake powder bottle to uniformly blend the special powders.
- 4. Dispense 1 full scoop of powder to a mixing pad or glass slab.
- 5. Dispense 2 drops of liquid into the mixing pad or glass slab. When dispensing liquid, hold the dropper vertically.
- 6. Incorporate powder into liquid in small increments. Total mixing time should not exceed 30 seconds
- 7. Apply the material, without delay, in a uniform layer.
- 8. Placement of mixed cement should be done immediately while mix is still glossy; approximately 1& 1/2 to 2 minutes working time. If cement becomes dull, it should be discarded and a new mixture prepared. If a longer working time is needed, mix the cement on a cold, dry slab.
- 9. Keep restoration free from saliva or water for 5 minutes.

Important Information:

Excess cement should be removed with a cotton roll immediately, before it sets, and should be cleaned from all surrounding areas such as gingiva and adjacent teeth before final set.

WARNING:

- 1. Medical devices should be kept out of reach of children.
- 2. DO NOT mix powder or liquid with any other glass ionomer components.

USE AND STORAGE:









Store away from sunlight

Observe storage temperatures

Protect from moisture

Shelf life, 3 years

EMERGENCY MEASURES:

Eye contact - Flush with copious amounts of water. Seek medical attention if irritation persists.

Ingestion - Dilute by drinking water.

Skin contact – Wash with soap and water.

CAUTION: Federal law restricts this device to sale by, or on the order of, a dentist (or dental professional).

International Orthodontic Services
12811 Capricorn St. Stafford, TX 77477 USA.
T: +1 (832) 342 9487 F: +1 (281) 340 4794
www.iosortho.com info@iosortho.com